

Presence of Different Bacterial Species in Thermal Sources and Novelty in their Industrial Enzyme Productions

Albayrak Seyda¹, Genc Berna², Adiguzel Gulsah³, Taskin Mesut^{1*} and Adiguzel Ahmet^{1*}

¹Department of Molecular Biology and Genetics, Faculty of Science, Ataturk University, Erzurum - 25240, Turkey. ²Department of Genetic and Bioengineering, Faculty of Engineering and Life Sciences, Gumushane University, Gumushane - 29100, Turkey. ³Department of Food Science and Technology, Faculty of Veterinary Science, Ataturk University, Erzurum - 25240, Turkey.

Abstract

In this study, one hundred and thirty isolates were isolated from water and sludge samples taken from hot springs located in different regions of Turkey. Among them, eleven isolates were chosen according to conventional (morphological, physiological and biochemical tests) and molecular methods (rep-PCR and 16S rRNA sequencing). These bacteria were then tested for their capability to produce valuable enzymes. As a result; species belonging to *Bacillus*, *Anoxybacillus*, *Aeribacillus*, *Enterococcus*, *Exiguobacterium* and *Paenibacillus* were identified. Test strains were found to have optimum reproductive potential at pH 5.0-9.0 and 15-65°C, usually at a concentration of 1.0-10.0% (w/v) NaCl. In addition, all thermo-tolerant bacteria were Gram, endospore (except *E. profundum*), catalase and oxidase (except *E. faecium* and *E. profundum*) positive, and rod-shaped (except *E. faecium*). It was observed that all isolates had a 99% similarity percentage as a result of 16S rRNA sequence analysis. All of the isolates were capable of producing industrially important enzymes moreover, eight of them could produce at least two of these enzymes. Test strains had high potential of industrial enzyme production, and the enzymes from these thermo-tolerant isolates will be widely used in biotechnological processes.

Keywords: Thermo-tolerant, isolation, 16S rRNA, rep PCR, industrial enzymes.

*Correspondence: mesut.taskin@atauni.edu.tr, adiguzel@atauni.edu.tr

(Received: 11 June 2019; accepted: 05 July 2019)

Citation: Albayrak Seyda, Genc Berna, Adiguzel Gulsah, Taskin Mesut and Adiguzel Ahmet, Presence of Different Bacterial Species in Thermal Sources and Novelty in their Industrial Enzyme Productions, *J Pure Appl Microbiol.*, 2019; **13**(3): doi: 10.22207/JPAM.13.3.

© The Author(s) 2019. **Open Access.** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

INTRODUCTION

The extraordinary diversity of life on earth has attracted researchers' attention from the past. Not only the diversity of living things, but also their distribution and function in different geographies is very attractive. However, an important part of the information on diversity is focused on the dissemination and protection of biodiversity on animals and plants. Most of the microbial diversity is still not discovered (>95%). Only 1% of the species diversity of prokaryotes in the world has been identified; therefore, little is known about prokaryotic diversity^{26,54,56}.

For each microorganism, there is the minimum temperature at which the breeding is not possible, the optimum temperature at which the breeding is fastest and the maximum temperature values at which the breeding is not possible. These three temperatures, which are called cardinal temperatures, are typical for each organism and vary significantly between species³⁴. Thermophilic bacteria which grow in hot springs, tropical soils, fertilizer heaps, manure forming fodder and garbage¹⁰ the temperature at which they can generally live optimally at 60°C. However, it is stated that very few thermophilic bacteria can live above 75°C²². In order to live and reproduce in extreme conditions, these organisms must adapt their metabolic and other cellular functions to these environments. The cell membrane of thermophilic is composed of saturated fatty acids. These fatty acids provide the cell with a hydrophobic environment and keep the cell tight and firm enough to survive at high temperatures. The cellular elements (cell membrane) and components (enzymes, proteins, nucleic acids, etc.) of thermophilic organisms are resistant to high temperatures. They are also resistant to denaturants such as extremely acidic and alkaline conditions and to proteolysis^{14,24}. The DNA of thermophiles contains a DNA Gyrase, which provides the reversible in the DNA to create positive supercoils. This increases the melting point of the DNA to at least the organism's maximum growth temperature. Thermophiles also tolerate high temperatures using increasing interactions such as electrostatic disulfide bridge and hydrophobic interactions using non-thermotolerant organisms²⁴.

Natural geothermal areas are found in tectonic active zone regions where earth crust movements in the world. Turkey is one of the richest countries in the world in the aspect of geothermal sources. However, the number of scientific studies on the determination of microorganism flora in thermal sources in different geographic regions of our country and determination of species with potential to be used in industry is very limited⁴¹. Until now, conventional methods based on phenotypic characters have been used in the diagnosis of thermophilic microorganisms in scientific studies conducted both in the world and in our country. The identification of microorganisms by morphological, physiological and biochemical tests has been insufficient to determine the differences between the desired microorganisms. Moreover, the conventional methods need long-term experiments, over labor, experienced and trained researchers, and the results are open to alternative interpretations, other known disadvantages of these methods³⁶. Therefore, in recent years, molecular techniques have been developed to be used in the diagnosis of microorganisms. Genetic profiles (16-23S rDNA, REP, ERIC and BOX-PCR) are commonly used methods³⁵.

Rai and Mukherjee⁴⁵ defined enzymes as green chemicals, specific biocatalysts that provide the realization and control of all reactions occurring in the living organism under suitable conditions, most of which are of protein structure and naturally produced only by living things. microbial enzymes which have applications in food industry, food and beverage production, cleaning and processing of garments in the textile industry, paper and detergent industries, medicine and medical field diagnosis of diseases continue to change with each passing day especially with biotechnological approaches and can be included in different application areas. The worldwide production volume of industrial enzymes, most of which have extracellular properties, was \$ 1 million in 1995; In 2000, approximately 60% of the industrial enzymes used in the world market were produced in Europe and 40% in the USA and Japan³⁰. The need for non-toxic and more effective methods has enabled the production of enzymes to scale up and penetrate into different markets thanks to the rapidly growing biotechnology.

Thermophilic organisms have a very important place due to their biotechnological potential. Their most important biotechnological features are; to produce enzymes that can catalyze biochemical reactions at much higher temperatures than normal organisms²³. These enzymes are thermophilic enzymes, and these enzymes are preferred in many industrial areas due to their pH changes and their stability to high temperatures. Thermostable enzymes play a role in the hydrolysis of protein, lipid and polysaccharide substrates in molecular biology⁷. Proteases from thermophilic enzymes are used in food industry, pharmacology, leather and detergent industry in a wide industrial area. enzymes such as lipase, esterase, amylase, cellulase and xylanase; used in detergent, sugar, textile and paper industries⁵⁵.

Thermostable enzymes are more advantageous than other enzymes, and both the industrially important and the thermostable enzyme producer, the introduction of new source microorganisms into the literature, will be able to respond to the needs of industry today and to shed light on possible future studies. The aim of the study was isolation and identification of new and industrially valuable bacterial strains from different geothermal resources of Turkey, presentation of their novel potential to produce industrially important enzymes and multi-enzyme productions.

MATERIALS AND METHODS

Purification of thermo-tolerant isolates

The water and sludge samples from Armutlu (Yalova), Germencik (Aydın), Yildizburnu (Izmir), Havza (Samsun), Guroymak (Bitlis) and Karakurt (Kırşehir) hot springs were collected and the samples were transferred to Nutrient Agar (NA) medium (peptone from meat 5.0 gL⁻¹; meat extract 3.0 gL⁻¹; agar 12.0 gL⁻¹) in order to incubate (Thermo Scientific Heratherm, the USA) at 55°C for 24 h. The growing colonies were spread on NA to obtain pure cultures and were eliminated by morphology differences in plates, as a first step. The pure, single and different colonies were stored in the Nutrient Broth (NB) with 15% glycerol content at -86°C for further studies³.

Gene amplification, cloning and genomic fingerprinting of isolates

For DNA extraction of each isolate, a

single colony from NA medium was selected and inoculated to NB at 55°C 150 rpm. After the incubation period, genomic DNA isolation was carried out according to Promega WizardR DNA purification kit and purified DNA was stored at +4°C until use⁴⁷. Test isolates were subjected to rep-PCR with the special primers of GTG₅ and BOX elements to obtain genomic fingerprinting^{4,6,20}. To obtain the PCR products, 50 ng of purified DNA was used as the template in 30 ml reaction mixture. Twenty-seven microliters of the reaction cocktail were prepared as follows: Specific Gitschier Buffer 5 mL, dimethyl sulfoxide 2.5 mL (100%, 20X), dNTPs (10 mM) 1.25 mL, bovine serum albumin 1.25 mL (20 gL⁻¹), primers (5 mM) 3.0 ml, Taq polymerase (250 U) 0.3 mL and water 13.7 mL. A negative control (no DNA) was included in each PCR assay. PCR reactions were performed with a 3x32-well ProFlex™ PCR System, using the following conditions for GTG₅-PCR: an initial denaturation at 94°C for 7 min; 36 cycles consisting of 94°C for 1 min and annealing at 53°C for 1 min with GTG₅ primer, extension at 65°C for 8 min; and a final polymerization at 65°C for 16 min before cooling at 4°C. For BOX-PCR; an initial denaturation at 95°C for 7 min; 36 cycles consisting of 94°C for 1 min and annealing at 53°C for 1 min with BOX primer, extension at 65°C for 8 min; and a final polymerization at 65°C for 16 min before cooling at 4°C.

The 16S rRNA of the isolates was amplified by Polymerase Chain Reaction (PCR) with the forward primer 27-F:(5' AGAGTTTGATYMTGGCTCAG3') and the reverse primer 1492-R:(5' GGTTACCTGTGACTT 3')⁵. The amplified fragments were cloned into *E. coli* JM101 strain with a vector system (pGEM-T, Promega, the UK) and the clones were sequenced (MacroGen, Amsterdam, the Netherlands). The results of 16S rRNA gene sequencing were analyzed using the GenBank (<http://blast.ncbi.nlm.nih.gov/blast.cgi>) and EzTaxon (<http://www.eztaxon.org/>) servers⁹. Considering the results of the study, a phylogenetic tree was formed via the neighbor-joining method by using the software package MEGA 4.0¹³.

The PCR products (27 mL) were mixed with 3 mL gel loading buffer (6X) and subjected to agarose (1.5% w/v) gel electrophoresis in Tris–Acetate–EDTA (TAE) buffer at 100 V for 110 min.

After separation of the amplification products by the gel, the fragments were stained with ethidium bromide solution (2 mL EtBr/100 mL 1X TAE buffer). The amplified DNA product was monitored using Gel Documentation System.

Conventional identification

The test isolates were subjected to conventional tests. The pH (Mettler Toledo) and temperature range for bacterial growth were tested between pH 3.0–11.0 and 15–65°C. The NaCl requirement for growth was also tested in NB medium containing 1.0, 2.0, 4.0, 6.0, 8.0, 10.0 and 12.0% (w/v) NaCl. Cell and colony morphology (Leica ICC50 HD light microscope), Gram and endospore staining, motility and the presence of catalase and oxidase reactions were also investigated by the method of Prescott⁴⁴ and Sari⁴⁷.

Preliminary experiments of industrial enzymes Amylase

Amylase assay was performed at 55°C for two days in a modified-medium which contained yeast extract (1 gL⁻¹), starch (5 gL⁻¹) and agar (15 gL⁻¹)⁵⁸. The amylase activity was screened by Lugol's solution. Clear and large zones were evaluated as positive, or as amylase production; however, no zone formation was evaluated as negative, or as no amylase production.

Cellulase

A medium containing carboxymethyl-cellulose (10 gL⁻¹), peptone (5 gL⁻¹), yeast (5 gL⁻¹), KH₂PO₄ (1 gL⁻¹), MgSO₄·7H₂O (0.2 gL⁻¹), NaCl (10 gL⁻¹) and agar (15 gL⁻¹) was used for cellulase assay. Incubation was carried out at 55°C for two days, and after the incubation, the plates were stained by Congo-Red Dye. The presence of clear zone was accepted as positive⁴⁷.

Lipase

Test isolates were inoculated on tributyrin agar medium which contained 1% (w/v) tributyrin (glycerol tributyrate) and incubated at 55°C. Formation of hydrolysis zone around the culture on plate was controlled for two days, and the strains with transparent and the highest zone formation (lipolytic activity) were determined as lipase producers¹².

Protease

Protease activity was conducted in medium which contained NB (8 gL⁻¹), Skimmed Milk (10 gL⁻¹) and agar (15 gL⁻¹) at 55°C for two

days. The plates were evaluated according to the zone formations, and the observation of a halo zone indicated positive protease activity⁴².

Xylanase

A solid medium with xylan (10 gL⁻¹), NaNO₃ (1.2 gL⁻¹), KH₂PO₄ (3 gL⁻¹), K₂HPO₄ (6 gL⁻¹), CaCl₂ (0.05 gL⁻¹), MgSO₄ (0.01 gL⁻¹), ZnSO₄ (0.001 gL⁻¹) and agar (15 gL⁻¹) was used to determine xylanase production. The plates were incubated at 55°C for four days and after the incubation, the plates were stained by Congo-Red Dye. The presence of clear zone was accepted as positive⁴⁷.

RESULTS AND DISCUSSION

Genomic fingerprinting of thermophiles

The first step of this study was the isolation of thermophilic bacteria from water and sludge samples of Armutlu (Yalova), Germencik (Aydın), Yildizburnu (Izmir), Havza (Samsun), Guroymak (Bitlis) and Karakurt (Kırşehir) hot springs. One hundred and thirty bacterial isolates were obtained however this number was firstly reduced by eliminating according to colony differences observed on plate surface. As conventional analysis alone was not sufficient for the identification of bacteria, rep-PCR [(GTG)₅-PCR and BOX-PCR] method was used which clearly showed the differences between microorganisms' genomic fingerprinting. (GTG)₅-PCR, gave the first information about isolates. It was observed that the isolates usually yielded up to sixteen polymorphic bands between 350-3500 bp and eight polymorphic bands between 700-2500 bp while EA 9 isolate gave no bands (Fig. 1). BOX-PCR, another genomic fingerprint method, was applied and it was determined that the test isolates were given two types of polymorphic bands, usually five bands between 300-1200 bp and two bands between 1200 - 4000 bp (Fig. 2). 16S rRNA region, which was evolutionarily conserved, was amplified using universal primers; to identify the isolates at species level. As a result of the analysis, eleven different species were detected, resulted in the fragments of 1500 bp. The sequences of cloned isolates were compared with the sequences of other bacteria in the GenBank and the nucleotides were analyzed using BLAST and EzTaxon (Table 1). The phylogenetic relationship between thermophilic bacteria was presented by phylogenetic tree (Fig. 3). As a result, the genus

Table 1. Molecular identifications of the isolates

Code	Name	Similarity ratio (%)	GenBank number
EA1	<i>Aeribacillus pallidus</i>	99.9	MH411153
EA2	<i>Anoxybacillus geothermalis</i>	99.7	MH411161
EA3	<i>Anoxybacillus mongoliensis</i>	99.3	MH411164
EA4	<i>Anoxybacillus rupiensis</i>	99.5	MH411201
EA5	<i>Bacillus albus</i>	99.8	MH411233
EA6	<i>Bacillus halodurans</i>	99.7	MH411241
EA7	<i>Bacillus licheniformis</i>	98.9	MH411598
EA8	<i>Bacillus paralicheniformis</i>	99.8	MH411597
EA9	<i>Enterococcus faecium</i>	99.9	MH411686
EA10	<i>Exiguobacterium profundum</i>	99.9	MH411710
EA11	<i>Paenibacillus dendritiformis</i>	99.7	MH411711

Bacillus with different species, *Enterococcus* and *Exiguobacterium* were determined. EA1 was similar to *Aeribacillus pallidus*, EA2 to *Anoxybacillus geothermalis*, EA3 to *Anoxybacillus mongoliensis*, EA5 to *Bacillus albus*, EA6 to *Bacillus halodurans*, EA7 to *Bacillus licheniformis*, EA8 to *Bacillus paralicheniformis*, EA9 to *Enterococcus faecium*, EA10 to *Exiguobacterium profundum* and EA11 to *Paenibacillus dendritiformis* species at a rate of 99%. Based on the knowledge that the similarity ratio of 16S rRNA sequence analysis among the strains of the same species should be more than 97%³⁴, it was concluded that the bacterial strains were similar to test strains.

Rep-PCR [(GTG)₅-PCR and BOX-PCR], which was molecular method used for this purpose, was a highly effective method used to reveal the diversity in the ecosystem, the phylogenetic relationship between the strains and to distinguish genetically close micro-organisms at species and sub-species level⁴. BOX-PCR and (GTG)₅-PCR methods were found to be very successful in demonstrating the genomic differences between strains in parallel with literature data^{3,4,6,58}.

Morphology, physiology and biochemical characteristics of thermophiles

Among eleven different bacterial isolates, all of isolates were Gram positive and motile, endospore producing (except EA10), catalase




Fig. 1. (GTG)₅ PCR band profiles of thermophilic isolates (M: DNA Marker, 1: EA 10, 2: EA 6, 3: EA 5, 4: EA 3, 5: EA 9, 6: EA 11, 7: EA 1, 8: EA 7, 9: EA 4, 10: EA 8, 11: EA 2, 12: Negative control)




Fig. 2. BOX-PCR band profiles of thermophilic isolates (M: DNA Marker, 1: EA 2, 2: EA 8, 3: EA 4, 4: EA 7, 5: EA 1, 6: EA 11, 7: EA 9, 8: EA 3, 9: EA 5, 10: EA 6, 11: EA 10, 12: Negative control)

and oxidase positive (except EA9 and EA10) and bacilli-shaped (EA9 was coccus-shaped) (Table 2). To determine the salt concentration for growth of the isolates, 1.0, 2.0, 4.0, 6.0, 8.0, 10.0 and 12% (w/v) NaCl were used (Table 2). As a result of spectrophotometric measurements made at

OD600, the test strains showed development in 1.0-10.0% salt concentration values; EA1 and EA4 was in 1.0-2.0%, EA3, EA7, EA8 and EA11 were in 2.0-4.0%, EA9 was in 2.0-6.0%, EA2 and EA6 were 2.0-8.0% and EA5 and EA10 were 2.0-10.0% range. In order to determine the pH ranges of test strains;

Table 2. Physicochemical requirements and conventional identifications of the isolates

Isolate code	Isolate name	Morphology	Gram	pH	Temp. (°C)	Salt (%)	Oxidase	Catalase	Endospore	Motility
EA1	<i>Aeribacillus pallidus</i>	Bacilli	+	7.0-9.0	35-55	1.0-2.0	+	+	+	+
EA2	<i>Anoxybacillus geothermalis</i>	Bacilli	+	5.0-9.0	35-55	2.0-8.0	+	+	+	+
EA3	<i>Anoxybacillus mongoliensis</i>	Bacilli	+	7.0-9.0	35-65	2.0-4.0	+	+	+	+
EA4	<i>Anoxybacillus rupiensis</i>	Bacilli	+	6.0-7.0	35-55	1.0-2.0	+	+	+	+
EA5	<i>Bacillus albus</i>	Bacilli	+	7.0-9.0	25-55	2.0-10.0	+	+	+	+
EA6	<i>Bacillus halodurans</i>	Bacilli	+	7.0-9.0	25-55	2.0-8.0	+	+	+	+
EA7	<i>Bacillus licheniformis</i>	Bacilli	+	7.0-9.0	35-65	2.0-4.0	+	+	+	+
EA8	<i>Bacillus paralicheniformis</i>	Bacilli	+	7.0-9.0	35-55	2.0-4.0	+	+	+	+
EA9	<i>Enterococcus faecium</i>	Coccus	+	5.0-9.0	15-55	2.0-6.0	-	+	*	+
EA10	<i>Exiguobacterium profundum</i>	Bacilli	+	5.0-9.0	15-55	2.0-10.0	-	+	-	+
EA11	<i>Paenibacillus dendritiformis</i>	Bacilli	+	6.0-7.0	35-55	2.0-4.0	+	+	+	+

*; endospore test was not applied.

microbial growth in pH 3.0, 5.0, 7.0, 9.0 and 11.0 were monitored; EA4 and EA11 were detected to develop in pH 6.0-7.0; EA1, EA3, EA5, EA6, EA7 and EA8 were in pH 7.0-9.0; EA2, EA9 and EA10 were in pH 5.0-9.0. The isolates were incubated at 15°C, 25°C, 35°C, 45°C, 55°C and 65°C to determine the temperatures at which they developed and the temperature values of the test strains were determined. Although the temperature ranges of thermophiles varied, EA9 and EA10 were determined that they could develop at 15-55°C, EA5 and EA6 were at 25-55°C, EA3 and EA7 were at 35-65°C, others were 35-55°C.

As a result of the morphological analysis of the isolates obtained in parallel with the literature data, all test strains were Gram and endospore (except *E. profundum*) positive, motile and had bacilli cell morphology except *E. faecium* (coccus). Growth ranges for pH, temperature and salt concentration were 5-9 pH, 15-65°C and 1-10%(w/v). Finally, all the isolates were catalase

and oxidase positive (except *E. faecium* and *E. profundum*). Yadav⁵⁷ carried out the isolation and identification of 150 thermophilic bacteria from hot springs in Nepal. The 16S rRNA fragments of the isolates were amplified with a size of about 1.5 kbs (1500 bp) and the microorganisms were belonging to the genus *Anoxybacillus*, *Aeribacillus*, and *Bacillus* at a ratio of ≥ 95 . The *bacilli* isolates were gram positive, endospore forming and their optimum growth temperature were between 55-65°C. However, in this study, the isolates were thermo-tolerant due to large temperate range, 15-65°C. Aanniz¹ isolated and identified 240 bacteria from different sources in Morocco. All isolates were Gram and endospore positive. When they examined the BOX-PCR and 16S rRNA sequence of the isolates, the dominant species was *Bacillus* (97.5%); 119 of them were *B. licheniformis* and 6 of the species belonging to *Aerobacillus*. Norashirene³⁸ obtained six thermophilic isolates from Malaysia. The optimum growth temperatures

Table 3. Industrial enzyme production profiles of the isolates and their first investments in thermophilic regions

Isolate code	Isolate name	A	C	L	P	X	Global	Turkey	Reference(s)
EA1	<i>Aeribacillus pallidus</i>	-	-	+	+	-	P	P	1, 12, 31, 57
EA2	<i>Anoxybacillus geothermalis</i>	+	-	+*	+	+*	P	*	19
EA3	<i>Anoxybacillus mongoliensis</i>	+	-	+*	+	+*	P	*	37
EA4	<i>Anoxybacillus rupiensis</i>	+	-	+*	+	-	P	P	16, 27, 58
EA5	<i>Bacillus albus</i> **	+*	-	+*	+*	-	*	*	33
EA6	<i>Bacillus halodurans</i>	-	+*	-	-	-	P	*	17, 25, 32, 46, 53
EA7	<i>Bacillus licheniformis</i>	+	-	+	+	+	P	P	1, 12, 58
EA8	<i>Bacillus paralicheniformis</i>	+*	-	+*	+*	+*	*	*	40
EA9	<i>Enterococcus faecium</i>	-	-	+*	+*	-	P	*	2, 39
EA10	<i>Exiguobacterium profundum</i>	-	-	+*	-	-	P	*	15
EA11	<i>Paenibacillus dendritiformis</i>	-	-	-	+*	-	*	*	49

(+):producer, (-): not producer; *: No study about related enzyme production ability/ No presence of bacteria at thermophilic regions; P:precence of the related bacteria in thermophilic regions**; <https://www.ncbi.nlm.nih.gov/genome/genomes/71466>, Abbreviations: A-Amylase; C-Cellulase; L-Lipase; P-Protease; X-Xylanase.

of these isolates were 55°C and pH was 7.5, and they were aerobic, catalase and oxidase positive. Poli⁴³ carried out the isolation of thermophilic bacteria by taking water and mud samples from the geothermal areas in Italy. These test strains were aerobic, endospore and Gram positive, motile, rod shaped and their optimum growth was at 65°C and pH 7.2. The salt concentrations in which the test strains could develop were tested at 1-10%(w/v), and the results were generally parallel

to the literature data. Tambekar⁵² revealed the presence of the strains that could reproduce in 8% (w/v) salt concentration, in parallel with our data. However, *B. halodurans* (EA6) was found to be able to withstand up to 8% (w/v) salt concentration in contrast to the literature⁵¹.

It was noticeable that the growth of *E. faecium*, a lactic acid bacterium at 15-55°C, considering that mesophilic bacteria generally did not show growth above 45°C. However, Orr³⁹




Fig. 3. Neighbor-Joining phylogenetic tree of thermophilic bacteria isolated from different hot springs of Turkey based on 16S rRNA gene analysis results. *Alicyclobacillus acidiphilus* used as an external group

found some members of the microorganisms contained in *Enterococci* were thermo-tolerant. vec⁵⁰ reported that, (GTG)₅-PCR method could be used effectively in the identification of the genus *Enterococcus* at species and sub-species level. Abdel-Rahman² identified *E. faecium* with 99% similarity ratio and as unable to utilize oxygen as an electron acceptor. They also concluded that; *E. faecium* could grow at 45-50°C and pH 9.6. Al-Mariri⁸ collected white cheese samples and identified thermophilic streptococci. As a result; *E. faecium* isolates were gram positive, non-sporulation and could grow at 4-6.5%(w/v) salt concentration while the salt concentration for this study was 2-6%(w/v). In another study; *E. faecium* was identified as catalase positive²¹ however Abdel-Rahman² and Jensen²⁸ identified as catalase negative. According to Schleifer and Kilpper-Bälz⁴⁸, some strains of *E. faecium* were motile. Crapart¹⁵ isolated lactic acid-producing thermo-tolerant bacterium, *E. profundum*, from hydrothermal vent and the growth conditions were parallel to our study with 12-49°C, 5.5-9.5 pH and 0-11%(w/v) salt concentration. This bacterium was characterized as motile, Gram and catalase positive, oxidase negative while Kasana and Pandey²⁹ also concluded as non-sporulation.

Enzyme production capacities of thermophiles

Enzyme production abilities of these eleven isolates were tested in enzyme specific solid media. As a result; all of them were producers of at least one enzyme which were industrially valuable (Table 3). EA2, EA3, EA4, EA5, EA7 and EA8 were amylase producers. The only cellulase producer isolate was EA6. Except EA6 and EA11, the rest of the isolates were positive in terms of lipase production. EA6 and EA10 gave negative results in protease growth media however, the other isolates were protease producers. EA2, EA3, EA7 and EA8 were xylanase producers.

A. pallidus, which was isolated from desert soil by Aanniz¹, could grow at 30-80°C, 0.5-10%(w/v) salt concentration and produce amylase and protease. Koc³¹ determined *A. pallidus* C196 and D642 as potential lipase producers due to use of olive oil as substrate. The isolation from different locations of Turkey resulted in identification of *A. pallidus* as amylase producer¹². With optimum growth conditions between 40-65°C, 0-3% (w/v) NaCl and 5.0-9.5

pH *A. geothermalis* could hydrolyze casein and starch due to protease and amylase activity, respectively¹⁹ while *A. geothermalis* in this study could also able to produce lipase and xylanase. Namsaraev³⁷ determined *A. mongoliensis*, whose optimum growth was at 35-75°C, 0-5% (w/v) NaCl concentration and 5.5-10.8 pH, as a thermophilic protease and amylase producer due to the hydrolysis of starch, casein and gelatin. *A. mongoliensis* in this study was also identified as lipase and xylanase producer. Dereková¹⁶ was isolated and identified *A. rupiensis*, which could survive between 35-67°C, 5.5-8.5 pH, for the first time with ability to degrade xylan, starch and casein. Jardine²⁷ was also proposed *A. rupiensis* as amylase, protease and cellulase producer. Yanmis⁵⁸ isolated *A. rupiensis* from geothermal regions of Turkey and it was declared as amylase producer. However, lipase production of *A. rupiensis* was determined in this study.

B. albus was deposited in NCBI under the project numbers of PRJNA509543, PRJNA326285 and PRJNA516150. Liu³³ identified the marine isolate as *B. albus* sp. nov., type strain N35-10-2T (=MCCC 1A02146T =KCTC 33710T =LMG 28875T). However, there is no publication about thermophilic source of the microorganism.

The lipolytic activity of *B. halodurans* was proved by many of the studies^{17,46,53} and Vargas⁵³ determined the optimum ranges at 25-55°C and 2.5-10% (w/v) NaCl. As a potential producer, amylase²⁵ and xylanase³² were other enzymes that *B. halodurans* produced. However, *B. halodurans* in this study was identified as cellulase producer. Aanniz¹ isolated *B. licheniformis*, which could survive at 30-75°C, 0.5-10% (w/v) salt content and 7-8.2 pH, had the potential to produce amylase, protease and cellulase. Archana and Satyanarayana¹¹ isolated cellulase-free xylanase from thermostable *B. licheniformis*. Baltaci¹² classified two of the isolates as *B. licheniformis* and they were also amylase, lipase, protease and cellulase producers. *B. licheniformis*, from thermal regions of Turkey⁵⁸, could also able to produce cellulase and amylase.

Although *B. paralicheniformis* was isolated from fermented foods^{18,40}, there was no record of thermophilic *B. paralicheniformis* and its enzyme production. *B. paralicheniformis* in this study could able to produce amylase, lipase, protease and

xylanase. *E. faecium* was firstly isolated by Orr³⁹ from hospital samples as thermo-tolerant however there was no report about enzyme production of thermo-tolerant *E. faecium*. However, lipase and protease production ability of *E. faecium* was determined in this study. Moderately thermophilic *E. profundum* was defined by Crapart¹⁵ but its industrial enzyme abilities never examined while lipase production of *E. profundum* discovered in this study. *P. dendritiformis*, which was included in the genus *Bacillus*, was first discovered in 1990s and reclassified as a separate genus⁴⁹. However, there is no record about *P. dendritiformis* from thermophilic sources while protease production was determined in this study.

In addition, the isolation locations of organisms were important during the isolation and identification of thermo-tolerant organisms (Table 3). In this study, the presence of *B. albus*, *B. paralicheniformis* and *P. dendritiformis* in thermophilic regions was introduced for the first time. The presence of *A. geothermalis*, *A. mongoliensis*, *B. albus*, *B. halodurans*, *B. paralicheniformis*, *E. faecium*, *E. profundum* and *P. dendritiformis* in the thermal sources of Turkey, has been detected for the first time. When amylase-producing isolates were screened, it was observed that most of the isolates were producers of amylase. However, among them, *B. albus* and *B. paralicheniformis* were firstly identified as thermo-tolerant amylase producers. Although thermophilic bacteria are frequently identified as cellulase producers, among the isolates in this study, only *B. halodurans* could produce cellulase and this was declared in this study. *A. pallidus*, *A. geothermalis*, *A. mongoliensis*, *A. rupiensis*, *B. albus*, *B. licheniformis*, *B. paralicheniformis*, *E. faecium* and *E. profundum* were thermo-tolerant bacteria with lipolytic activity. However, the lipase production abilities of the isolates, except *A. pallidus* and *B. licheniformis*, were not determined before, until this study. While all isolates, except *B. halodurans* and *E. profundum*, were able to produce protease, the proteolytic activities of thermo-tolerant *B. albus*, *B. paralicheniformis*, *E. faecium* and *P. dendritiformis* were determined for the first time. *A. geothermalis*, *A. mongoliensis*, *B. licheniformis* and *B. paralicheniformis* were able to produce xylanase; however, *A. geothermalis*,

A. mongoliensis and *B. paralicheniformis* were firstly determined as thermo-tolerant xylanase producers

When the multienzyme production capabilities of the isolates were examined, it was observed that all the isolates, except *B. halodurans*, *E. profundum* and *P. dendritiformis*, were able to produce more than one enzyme. In a study, in order to determine the strains which have the potential to produce industrially important enzymes, Yadav⁵⁷ observed that 135 of 150 thermophilic isolates have the potential to produce at least one extracellular hydrolytic enzyme.

CONCLUSION

As a conclusion, this research contained remarkable points in terms of presenting different thermo-tolerant species in the World and Turkey, microbial flora of thermal regions in Turkey and unknown abilities of these thermo-tolerant species to produce industrially valuable enzymes. The first isolation of *B. albus*, *B. paralicheniformis* and *P. dendritiformis* from thermal regions in the World was carried out. *A. geothermalis*, *A. mongoliensis*, *B. albus*, *B. halodurans*, *B. paralicheniformis*, *E. faecium*, *E. profundum* and *P. dendritiformis* were firstly isolated from hot springs of Turkey. *A. geothermalis*, *A. mongoliensis*, *B. albus*, *B. halodurans*, *B. paralicheniformis*, *E. faecium*, *E. profundum* and *P. dendritiformis* were identified as producers of some of industrial enzymes, for the first time. Eight different species with the potential to produce multi-enzyme were identified. Since thermostable enzymes play an important role in industrial and biotechnological processes, the enzyme/multi-enzyme production potentials of the isolates identified in this study has been one of the another notable results.

ACKNOWLEDGEMENTS

None

AUTHORS' CONTRIBUTION

AS and GB designed and carried out the study, GB and AG participated in design to draft the manuscript, TM and AA dealt with financial supports and funding. All the authors read and approved the final manuscript.

FUNDING

The authors acknowledge the support of Ataturk University, Turkey for this work. This work was supported by the Research Development Centre of Ataturk University (Grant numbers were 2015/54 and FAD-2018-6352).

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ETHICS STATEMENT

All text, data, figures/tables or other illustrations presented in the manuscript are completely original and does not contain or include material taken from other copyrighted sources. This article does not contain any studies about human or animal objects.

REFERENCES

- Aanniz T., Ouadghiri M., Melloul M., Swings J., Elfahime E., Ibjibij J., Ismaili M. and Amar M., Thermophilic bacteria in Moroccan hot springs, salt marshes and desert soils. *Brazilian Journal of Microbiology*, 2015; **46**: 443-453.
- Abdel-Rahman M.A., Tashiro Y., Zendo T., Sakai K. and Sonomoto K., Enterococcus faecium QU 50: a novel thermophilic lactic acid bacterium for high-yield l-lactic acid production from xylose. *FEMS Microbiology Letters*, 2015; **362**: 1-7.
- Adiguzel A., Molecular characterization of thermophilic bacteria isolated from water samples taken from various thermal plants. PhD, Atat rk University, Graduate School at Natural and Applied Sciences, Erzurum, Turkey, 2006; **130**.
- Adiguzel G., Bozoglu C., Yanmis D., Gormez A., Gulluce M. and Adiguzel A., Phenotyping and genotyping characterization of Salmonella strains isolated from retail beef in Erzurum, Turkey. *Journal of Pure and Applied Microbiology*, 2012; **6**: 1581-1589.
- Adiguzel G., Faiz O., Sisecioglu M., Sari B., Baltaci O., Akbulut S., Genc B and Adiguzel A., A novel endo- β -1,4-xylanase from Pediococcus acidilactici GC25; purification, characterization and application in clarification of fruit juices. *International Journal of Biological Macromolecules*, 2019; **129**: 571-578.
- Adiguzel G., Gulluce M., Bozoglu C., Yanmis D., Gormez A., Atasever M. and Adiguzel A., Molecular characterization of Escherichia coli O157: H7 from retail beef in Erzurum, Turkey. *Journal of Pure and Applied Microbiology*, 2012; **6**: 1033-1041.
- Aguilar A., Extremophile research in the European Union: from fundamental aspects to industrial expectations. *FEMS Microbiology Reviews*, 1996; **18**: 89-92.
- Al-Mariri A., Younes A.A. and Sharabi N-E., Characterization of thermophilic streptococci isolated from rustic white cheese. *The Journal of General and Applied Microbiology*, 2013; **59**: 97-103.
- Altschul S.F. and Gish W., [27] Local alignment statistics, in *Methods in Enzymology*. Academic Press, 1996; 460-480.
- Aragno M. and Kristjansson J., Aerobic, chemolithoautotrophic, thermophilic bacteria. *Thermophilic Bacteria*, 1992; 77-103.
- Archana A. and Satyanarayana T., Xylanase production by thermophilic Bacillus licheniformis A99 in solid-state fermentation. *Enzyme and Microbial Technology*, 1997; **21**: 12-17.
- Baltaci M.O., Genc B., Arslan S., Adiguzel G. and Adiguzel A., Isolation and Characterization of Thermophilic Bacteria from Geothermal Areas in Turkey and Preliminary Research on Biotechnologically Important Enzyme Production. *Geomicrobiol. J.*, 2017; **34**: 53-62.
- Benson G., Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.* 1999; **27**: 573.
- Bjarni A. and Magn'??s MKn, Properties of Extremophilic Enzymes and Their Importance in Food Science and Technology, in *Handbook of Food Enzymology*. CRC Press, 2002.
- Crapart S., Fardeau M-L., Cayol J-L., Thomas P., Sery C., Ollivier B. and Combet-Blanc Y., Exiguobacterium profundum sp. nov., a moderately thermophilic, lactic acid-producing bacterium isolated from a deep-sea hydro-thermal vent. *International Journal of Systematic and Evolutionary Microbiology*, 2007; **57**: 287-292.
- Derekova A., Sjrholm C., Mandeva R. and Kambourova M., Anoxybacillus rupiensis sp. nov., a novel thermophilic bacterium isolated from Rupi basin (Bulgaria). *Extremophiles*, 2007; **11**: 577-583.
- Dua A. and Gupta R., Functional characterization of hormone sensitive-like lipase from Bacillus halodurans: synthesis and recovery of pNP-laurate with high yields. *Extremophiles*, 2017; **21**: 871-889.
- Dunlap C.A., Kwon S-W., Rooney A.P. and Kim S-J., Bacillus paralicheniformis sp. nov., isolated from fermented soybean paste. *International Journal of Systematic and Evolutionary Microbiology*. 2015; **65**: 3487-3492.
- Filippidou S., Jaussi M., Junier T., Wunderlin T., Jeanneret N., Palmieri F., Palmieri I., Roussel-Delif L., Vieth-Hillebrand A and Vetter A, Anoxybacillus-geothermalis sp. nov., a facultatively anaerobic, endospore-forming bacterium isolated from mineral deposits in a geothermal station. *International Journal of Systematic and Evolutionary Microbiology*, 2016; **66**: 2944-2951.
- Gevers D., Huys G. and Swings J., Applicability of rep-PCR fingerprinting for identification of Lactobacillus species. *FEMS Microbiology Letters*, 2001; **205**: 31-36.
- Ghattargi V.C., Nimonkar Y.S., Burse S.A., Davray D., Kumbhare S.V., Shetty S.A., Gaikwad M.A., Suryavanshi M.V., Doijad S.P. and Utage B., Genomic and physiological analyses of an indigenous strain, Enterococcus faecium 17OM39. *Functional & Integrative Genomics*, 2018; **18**: 385-399.
- Guagliardi A., Martino M., Iaccarino I., Rosa M.D., Rossi

- M. and Bartolucci S., Purification and characterization of the alcohol dehydrogenase from a novel strain of *Bacillus stearothermophilus* growing at 70°C. *The International Journal of Biochemistry & Cell Biology*, 1996; **28**: 239-246.
23. Gugliandolo C., Maugeri T.L., Caccamo D. and Stackebrandt E., *Bacillus aeolius* sp. nov. a novel thermophilic, halophilic marine *Bacillus* species from Eolian Islands (Italy). *Syst. Appl. Microbiol.*, 2003; **26**: 172-176.
 24. Haki G. and Rakshit S., Developments in industrially important thermostable enzymes: a review. *Bioresource Technology*, 2003; **89**: 17-34.
 25. Hashim S.O., Delgado O.D., Martinez M.A., Kaul R-H., Mula F.J. and Mattiasson B., Alkaline active maltohexaose-forming α -amylase from *Bacillus halodurans* LBK 34. *Enzyme and Microbial Technology*, 2005; **36**: 139-146.
 26. Hugenholtz P. and Pace N.R., Identifying microbial diversity in the natural environment: a molecular phylogenetic approach. *Trends in biotechnology*, 1996; **14**: 190-197.
 27. Jardine J., Stoychev S., Mavumengwana V. and Ubomba-Jaswa E., Screening of potential bioremediation enzymes from hot spring bacteria using conventional plate assays and liquid chromatography-Tandem mass spectrometry (LC-MS/MS). *Journal of Environmental Management*. 2018; **223**: 787-796.
 28. Jensen M., Arda Y. and Vogensen F.K., Isolation of cultivable thermophilic lactic acid bacteria from cheeses made with mesophilic starter and molecular comparison with dairy related *Lactobacillus helveticus* strains. *Letters in Applied Microbiology*. 2009; **49**: 396-402.
 29. Kasana R.C. and Pandey C., Exiguobacterium: an overview of a versatile genus with potential in industry and agriculture. *Critical Reviews in Biotechnology*, 2018; **38**: 141-156.
 30. Kirk O., Borchert T.V. and Fuglsang C.C., Industrial enzyme applications. *Current Opinion in Biotechnology*, 2002; **13**: 345-351.
 31. Koc M., Cokmus C. and Cihan A.C., The genotypic diversity and lipase production of some thermophilic bacilli from different genera. *Brazilian Journal of Microbiology*, 2015; **46**: 1065-1076.
 32. Lin X-Q., Han S-Y., Zhang N, Hu H., Zheng S-P, Ye Y-R. and Lin Y., Bleach boosting effect of xylanase A from *Bacillus halodurans* C-125 in ECF bleaching of wheat straw pulp. *Enzyme and Microbial Technology*, 2013; **52**: 91-98.
 33. Liu Y., Du J., Lai Q., Zeng R., Ye D., Xu J. and Shao Z., Proposal of nine novel species of the *Bacillus cereus* group. *International Journal of Systematic and Evolutionary Microbiology*, 2017; **67**: 2499-2508.
 34. Madigan M.T., Martinko J.M. and Brock T.D., Brock biology of microorganisms. Pearson Prentice Hall Upper Saddle River, NJ., 2006.
 35. Manceau C. and Horvais A., Assessment of genetic diversity among strains of *Pseudomonas syringae* by PCR-restriction fragment length polymorphism analysis of rRNA operons with special emphasis on *P. syringae* pv. tomato. *Applied and Environmental Microbiology*, 1997; **63**: 498-505.
 36. Miller SA, T. R. Joaquim Diagnostik techniques for plant pathogens, *Biotechnology in Plant Disease Control*, 1993.
 37. Namsaraev Z., Babasanova O., Dunaevsky Y., Akimov V., Barkhutova D., Gorlenko V. and Namsaraev B., *Anoxybacillus mongoliensis* sp. nov., a novel thermophilic proteinase producing bacterium isolated from alkaline hot spring, central Mongolia. *Microbiology*, 2010; **79**: 491-499.
 38. Norashirene M., Zakiah Y., Nurdiana S., Hilwani I.N., Khairiyah M.S. and Arif M.M., Identification of Cellulose-Hydrolytic Thermophiles Isolated from Sg. Klah Hot Spring Based On 16S rDNA Gene Sequence. World Academy of Science, Engineering and Technology, *International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering*, 2014; **8**: 1041-1044.
 39. Orr K., Holliday M., Jones A., Robson I. and Perry J., Survival of enterococci during hospital laundry processing. *Journal of Hospital Infection*, 2002; **50**: 133-139.
 40. Othoum G., Bougouffa S., Razali R., Bokhari A., Alamoudi S., Antunes A., Gao X., Hoehndorf R., Arold S.T. and Gojobori T., In silico exploration of Red Sea *Bacillus* genomes for natural product biosynthetic gene clusters. *BMC Genomics*, 2018; **19**: 382.
 41. Ozel N. General Outlook on Geothermal Resources in GAP Region and Thermal Springs of Guclukonak (SIRNAK) District Geothermal Energy. Available from: www.emo.org.tr/ekler/ecbba75f70a1293_ek.doc, 2000.
 42. Panda M.K., Sahu M.K. and Tayung K., Isolation and characterization of a thermophilic *Bacillus* sp. with protease activity isolated from hot spring of Tarabalo, Odisha, India. *Iranian Journal of Microbiology*, 2013; **5**: 159.
 43. Poli A., Romano I., Cordella P., Orlando P., Nicolaus B. and Berrini C.C., *Anoxybacillus thermarum* sp. nov., a novel thermophilic bacterium isolated from thermal mud in Euganean hot springs, Abano Terme, Italy. *Extremophiles*, 2009; **13**: 867-874.
 44. Prescott L.M., Harley J.P. and Klein D.A., Laboratory exercises in microbiology. McGraw-Hill Companies, 2002.
 45. Rai S.K. and Mukherjee A.K., Statistical optimization of production, purification and industrial application of a laundry detergent and organic solvent-stable subtilisin-like serine protease (Alzwpiprase) from *Bacillus subtilis* DM-04. *Biochemical Engineering Journal*, 2010; **48**: 173-180.
 46. Ramchuran S.O., Vargas V.A., Hatti-Kaul R. and Karlsson E.N., Production of a lipolytic enzyme originating from *Bacillus halodurans* LBB2 in the methylotrophic yeast *Pichia pastoris*. *Applied Microbiology and Biotechnology*, 2006; **71**: 463-472.
 47. Sari B., Faiz O., Genc B., Sisecioglu M., Adiguzel A. and Adiguzel G., New xylanolytic enzyme from *Geobacillus galactosidasius* BS61 from a geothermal resource in Turkey. *International Journal of Biological Macromolecules*, 2018; **119**: 1017-1026.
 48. Schleifer K.H. and Kilpper-Bälz R., Transfer of

- Streptococcus faecalis* and *Streptococcus faecium* to the genus *Enterococcus* nom. rev. as *Enterococcus faecalis* comb. nov. and *Enterococcus faecium* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, 1984; **34**: 31-34.
49. Sirota-Madi A., Olender T., Helman Y., Brainis I., Finkelshtein A., Roth D., Hagai E., Leshkowitz D., Brodsky L., Galatenko V., Nikolaev V., Gutnick D.L., Lancet D. and Ben-Jacob E., Genome sequence of the pattern-forming social bacterium *Paenibacillus dendritiformis* C454 chiral morphotype. *Journal of Bacteriology*, 2012; **194**: 2127-2128.
50. Vec P., Vancanneyt M., Seman M., Snauwaert C., Lefebvre K., Sedláček I and Swings J, Evaluation of (GTG) 5-PCR for identification of *Enterococcus* spp. *FEMS Microbiology Letters*, 2005; **247**: 59-63.
51. Takami H., Akiba T. and Horikoshi K., Production of extremely thermostable alkaline protease from *Bacillus* sp. no. AH-101. *Applied Microbiology and Biotechnology*, 1989; **30**: 120-124.
52. Tambekar D., Tambekar S., Jadhav A. and Babhulkar B., Isolation and partial characterization of protease from *Bacillus halodurans* (AJ302709) from alkaline Lonar lake. *International Journal of Pharmaceutical Sciences and Research*, 2016; **7**: 4546.
53. Vargas V.A., Delgado O.D., Hatti-Kaul R. and Mattiasson B., Lipase-producing microorganisms from a Kenyan alkaline soda lake. *Biotechnology Letters*, 2004; **26**: 81-86.
54. Ward D.M., Weller R. and Bateson M.M., 16S rRNA sequences reveal numerous uncultured microorganisms in a natural community. *Nature*, 1990; **345**: 63.
55. Wery N., Cambon-Bonavita M-A., Lesongeur F. and Barbier G., Diversity of anaerobic heterotrophic thermophiles isolated from deep-sea hydrothermal vents of the Mid-Atlantic Ridge. *FEMS Microbiology Ecology*, 2002; **41**: 105-114.
56. Whitman W.B., Coleman D.C. and Wiebe W.J., Prokaryotes: the unseen majority. *Proceedings of the National Academy of Sciences*, 1998; **95**: 6578-6583.
57. Yadav P., Korpole S., Prasad G.S., Sahni G., Maharjan J., Sreerama L. and Bhattarai T., Morphological, enzymatic screening, and phylogenetic analysis of thermophilic bacilli isolated from five hot springs of Myagdi, Nepal. *Journal of Applied Biology & Biotechnology Vol.*, 2018; **6**: 1-8.
58. Yanmis D., Baltaci M.O., Gulluce M. and Adiguzel A., Identification of thermophilic strains from geothermal areas in Turkey by using conventional and molecular techniques. *Res. J. Biotechnol.*, 2015; **10**: 39-45.