

<https://doi.org/10.17221/9/2020-VETMED>

The impact of dietary tarragon (*Artemisia dracunculus*) on serum apelin, brain-derived neurotrophic factor, cardiac troponin concentrations and histopathology of liver tissue in laying hens housed at different stocking densities

BULENT BAYRAKTAR^{1*}, EMRE TEKCE², HACER KAYA³, MUSA KARAALP⁴, EYLUL TURUNC⁵

¹Faculty of Health Sciences, Bayburt University, Bayburt, Turkey

²Faculty of Applied Sciences, Bayburt University, Bayburt, Turkey

³Gümüşhane University Şiran Vocational School, Department of Veterinary Medicine, Şiran/ Gümüşhane, Turkey

⁴Gümüşhane University Kelkit Aydın Doğan Vocational School, Department of Veterinary Medicine, Kelkit/Gümüşhane, Turkey

⁵Department of Pathology Science, Atatürk University Erzurum, Turkey

*Corresponding author: bulenttbayraktar@gmail.com

Citation: Bayraktar B, Tekce E, Kaya H, Karaalp M, Turunc E (2020): The impact of dietary tarragon (*Artemisia dracunculus*) on serum apelin, brain-derived neurotrophic factor, cardiac troponin concentrations and histopathology of liver tissue in laying hens housed at different stocking densities. Vet Med-Czech 65, 269–279.

Abstract: Due to its association with several other stress factors (poultry house gases, inadequate ventilation, heat, cold and poor hygiene), the high stocking density is a major stress factor that adversely affects the health and performance of poultry and the quality of the poultry products. Therefore, this experimental study was aimed at analysing the impact of different doses of dietary tarragon (*Artemisia dracunculus*) on the serum apelin, plasma brain-derived neurotrophic factor (p-BDNF), and cardiac troponin I (cTnI) concentrations, and the correlation between these indicators in laying hens housed at different stocking densities. The aim of this study is to investigate the effects of adding tarragon in different ratios to laying hen rations in the 2nd ovulation period on the cTnI, apelin, and BDNF hormone concentrations and the liver histopathology. The experiment was carried out over a period of eight weeks, with 192 Lohman Brown commercial hybrids at 50 weeks of age. Eight groups (four replicates each), composed of laying hens of equal body weight, which were housed at stocking densities of 580 cm²/hen and 810 cm²/hen and received 0, 1, 5 and 10 mmol/kg of tarragon (*Artemisia dracunculus*) in the feed, were established. At the end of the trial, 96 of the housed egg-laying hens (3 birds in each subgroup, a total of 12 birds in each group) were randomly selected and blood samples were taken from the *vena subcutanea ulnaris*. The samples collected were analysed for the apelin, p-BDNF, and cTnI contents. The analysis results demonstrated that tarragon supplementation had no effect on the serum apelin, p-BDNF and cTnI concentrations ($P > 0.05$). The Sub-Groups ST1, ST1.2, and ST6 presented with severe hyperaemia of the sinusoidal, portal and acinar blood vessels, whilst the hyperaemia of these blood vessels was moderate in Sub-Group ST12. Apelin, BDNF, and cTnI can act as protective factors against negative consequences of stress (e.g., stocking density or heat stress).

Keywords: diet; diet supplementation; apelin; BDNF; cardiac troponin I (cTnI)

Poultry farming has a strategic importance in the livestock sector because of its role in meeting the demand for animal protein (meat, eggs) brought about by the continuous increase of the world's human population (Bayraktar and Tekce 2019). Commercial egg production is a major livestock sector, which faces multiple risks throughout the production and management processes, starting from the establishment of the farm, and extending to aspects such as disease prevention, animal nutrition and animal husbandry. Laying hen farms aim to maximise the profit per animal unit. A higher number of hens per square meter reduces the production costs, yet an excessive stocking density has a negative impact on the performance. The welfare of laying hens kept in conventional battery cages has been well scrutinised. Although conventional battery cages have been perceived as the most profitable housing system for layers, they are not considered to be welfare-friendly (Appleby 2004; Appleby et al. 1993; Craig and Swanson 1994). A high stocking density, resulting from the housing of a greater number of hens per square meter of usable area, is a major stress factor which causes overcrowding, behavioural disorders, reduced access to feed and water, increased rates of wounding and diseases, and eventually, poor welfare (Appleby 2004; Kang et al. 2016). A high stocking density has also been reported to cause dermatitis (Sorensen et al. 2000; Matkovic et al. 2019), abdominal wounds, skin lesions (Weimer et al. 2019) and thoracic oedema (Meseret 2016). The cage stocking density is classified under three categories as follows: high (384 cm²/hen), medium (464 cm²/hen), and low (580 cm²/hen) (McGlone 2010; Nicol et al. 2017). A high stocking density is reported to cause sudden behavioural changes and sudden death syndrome as well as several direct and indirect effects on the physiology and anatomy of the animals (Bessei 2006). The hypothalamic-pituitary-adrenal (HPA) axis, comprised of the hypothalamus (H), pituitary gland (P) and adrenal glands (A), regulates several physiological processes in the body, including digestion, immunity and energy storage among others. In response to a stressful stimulus, the hypothalamus releases vasopressin and a corticotropin-releasing hormone (CRH), which initiate the adrenocorticotrophic hormone (ACTH) synthesis in the pituitary gland. Furthermore, the release of stress hormones, referred to as corticotropins, as well as cortisol, and glucocorticoids, such as

corticosterone (CORT), used as stress indicators in animals, activate the HPA axis. Plasma cortisol or corticosterone is the primary glucocorticoid, the level of which is used as an indicator of stress and the endocrine response of the HPA axis to the stress (Charmandari et al. 2005).

In poultry species, egg production is an energy-intensive process. Excessive energy consumption reduces the laying capacity of the hens. Similar to the case in mammals, alterations in the lipid synthesis and metabolism cause granulosa cell apoptosis, altered immune functions and hormonal synthesis in chickens, and, thereby, place ovarian functions under risk (Walzem and Chen 2014). Apelin, which was first isolated from bovine stomach extracts and is known to originate from a 77 amino acid-precursor (preproapelin), is an adipokine involved in energy regulation (Tatemoto et al. 2001). Apelin, which is the endogenous ligand of the apelin (APJ) receptor, occurs in various isoforms, such as apelin-12, -13, -17, and -36. In terms of biological activity, isoform apelin-13 is 8 times stronger than apelin-17 and 60 times stronger than apelin-36 (Tatemoto et al. 1998). Therefore, the majority of recent research has focused on apelin-13, which has a higher biological activity than the other apelin isoforms and contains N-terminal pyroglutamate residues (Beltowski 2006). Cardiac troponin I (cTnI), which is an indicator of stress-related cardiac arrhythmia and myocardial damage, has found common use in the past decade as a highly specific cardiac marker (Adams et al. 1994; Wu et al. 1996; Brown and Bertolet 1997). Cardiac troponins are proteins which control the calcium-mediated interaction between actin and myosin and allow contraction at the sarcomere level. Troponins are major structural proteins, which, together with tropomyosin, have an important role in regulating the contraction of the skeletal and cardiac muscles (Hi et al. 2019). The troponin complex, which consists of three protein subunits, namely, troponin I, troponin T (cTnT), and troponin C (cTnC), and is located on the thin filament of the contractile apparatus, plays a significant role in the transmission of the intracellular calcium signal into the actin-myosin interaction. The amount of cardiac troponin, which passes into the blood circulation, depends on the type, duration and severity of the myocyte damage. The monitoring of the blood cTnI and cTnT concentrations is a biochemical method used to determine acute coronary syndromes

<https://doi.org/10.17221/9/2020-VETMED>

(Alpert et al. 2000). In the event of myocardial damage, troponins pass from the myocytes into the blood circulation. Depending on the severity of the myocardial damage, blood troponin levels elevate within 1–4 h on average, and are detectable for a period of 7–14 days, which makes it possible to assess and quantify the damage to the myocardium (Apple 1999).

As a growth mediator responsible for neurogenesis, the brain-derived neurotrophic factor (BDNF) regulates the growth, survival, and differentiation of the neurons and prevents ischaemia-induced cell death, and thereby, maintains the continuity of the cell activity and repair (Kertes et al. 2017). BDNF induces and controls the generation of neurons from stem cells (neurogenesis) (Zigova et al. 1998; Benrasis et al. 2001). Not only does it show a neuroprotective effect in the presence of stress (Spina et al. 1992), BDNF also has a fundamental role in energy homeostasis (Bothwell 1995). Stress is a neuronal damage factor, which reduces the BDNF levels (Lee et al. 2008; Fuchikami et al. 2009) and triggers degenerative cellular processes in the limbic system (Adlard and Cotman 2004; McEwen 2006). Brain lateralisation is reported to be of relevance to the assessment of animal welfare (Rogers et al. 2010).

Artemisia dracuncululus L., known as estragon or tarragon, is a small, shrubby, perennial plant, which belongs to the family Asteraceae and is native to Anatolia (Ceylan 1996; Kordali et al. 2005). Tarragon has a wide range of pharmacological activity, including antioxidant, antimicrobial, carminative, digestive, anti-inflammatory, antipyretic, antiseptic, antispasmodic, antiparasitic, anthelmintic, and fungicidal effects (Volak and Stodola 1998; Duke 2002; Hassanzadeh et al. 2016). Furthermore, this plant has also been reported to show effects on cerebral and gastrointestinal functions (Aglarova et al. 2008). There is no legal restriction on the use of tarragon as its commercially available form is classified as a non-toxic essential oil (Voitkevich 1999). Dried tarragon contains 24% protein, 45% carbohydrates, 7% fat and 7% fibre (Attokaran 2011). Tarragon also contains an essential oil (0.4–0.8%), bitter substance and tannin. Tarragon leaves are reported to contain 4% of an aromatic essential oil, the composition of which includes various substances such as camphor, artemisia ketone, and cineole (Mansuroglu and Gurel 2001).

Our hypothesis was that the addition of tarragon to the diet of laying hens would elicit a response in the form of serum and plasma adipokines (Apelin, BDNF, and cTnI).

The study was aimed at the investigation of the impact of different doses of dietary tarragon (*Artemisia dracuncululus*) on the serum apelin, plasma brain-derived neurotrophic factor (p-BDNF), cardiac troponin I (cTnI) concentrations and the histopathology of the liver tissue and the correlation between these variables in the egg-laying hens housed at different stocking densities.

MATERIAL AND METHODS

Animals, experimental design and feed

The study was conducted at the Kelkit Organic Agriculture Research and Application Centre of Gümüşhane University, and the animal material comprised 50-week-old, 192 hybrid commercial layers of the Lohmann Brown strain. After a 2 week-acclimatisation period, the study was carried out for a period of 8 weeks. Replicate groups composed of five and seven hens were established for the normal and high stocking densities, respectively. The size of the cages, in which the replicate groups were housed, was 90 × 45 × 35 cm, and the floor area per bird was 810 cm² at the normal stocking density and 580 cm² at the high stocking density. The feed ration provided to the laying hens was supplemented with ground homogenous tarragon at doses of 0, 1.2, 6 and 12 g/kg in the Groups T₀, T_{1.2}, T₆ and T₁₂, respectively. The study was conducted in compliance with the animal welfare rules and principles, pursuant to the approval, dated May 10, 2017 and numbered 2017/1-07, of the Local Ethics Board for Animal Experiments of Gümüşhane University.

Temperature, humidity and lighting of the poultry house

The laying hens were maintained on a 16:8 light-dark cycle and were provided with water and feed *ad libitum*. The temperature and humidity inside the poultry house were adjusted according to the needs of the animals. The composition of the commercial feed provided to the birds is presented in Table 1. The feed was analysed in accordance

Table 1. Basal diet ration nutrient content and analysis (g/kg)

Ingredients	%
Corn	35.00
Triticale	17.50
Wheat	7.50
Soybean meal (34%)	11.46
Sunflower meal (33%)	10.36
Hazelnut (42%)	3.50
Corn gluten meal (60%)	2.50
Vegetable oil	0.83
Limestone	9.55
Dicalcium phosphate (18%)	0.72
NaCl	0.32
Premix ¹	0.25
L-lysine HCl	0.15
Naturabind-S	0.12
Chemical analysis of feed (%)	
Crude matter	89.22
Crude protein	16.75
Ether extract	4.66
Crude fibre	5.29
Crude ash	13.17
Starch	33.99
Calculated contents of feed	
Metabolizable energy (kcal/kg)	2 735
Methionine (%)	0.37
Methionine + cystine (%)	0.67
Lysine (%)	0.75
Linoleic acid (%)	2.21
Ca (%)	3.90
Available P (%)	0.35
Na (%)	0.15

¹Premix provided per kilogram of diet: vitamin A, 10 000 IU; vitamin D3, 2 400 IU; vitamin E, 30 mg; vitamin K3, 2.5 mg; vitamin B1, 3 mg; vitamin B2, 7 mg; vitamin B6, 4 mg; vitamin B12, 0.02 mg; niacin, 40 mg; Ca-D-pantothenate, 8 mg; folic acid, 1 mg; D-biotin, 0.1 mg; vitamin C, 50 mg; choline chloride, 125 mg; Mn, 80 mg; Fe, 60 mg; Zn, 60 mg; Cu, 5 mg; Co, 0.10 mg; Se, 0.15 mg

with the methods adopted by the Association of Official Analytical Chemists (AOAC). During the acclimatisation period, a 16:8 light-dark cycle (60 W) was maintained and the temperature was adjusted to 22 °C.

Content of the tarragon (*Artemisia dracunculus*) leaves

The tarragon leaves added to the feed of the animals were obtained from farmers, who grew tarragon in the Yedigözüler village of the Bayburt province. After washing and removing the soil, plants and also non-usable parts of the herb, it was placed on a clean floor, and dried under the appropriate room temperature. The dried tarragon samples were powdered in a mill, and were added to the experimental diets (Hosseinzadeh and Moghaddam 2014). *Artemisia dracunculus* also has important compounds attributed to it, such as: methyl chavicol, ocimene, myrcene, camphor, camphene, *p*-anisic acid, limonene, linalool, *p*-methoxy cinnamic aldehyde, flavonoid, coumarin and minerals (Gulpinar 2012). The total antioxidant level of the tarragon plant used in this study was determined by the colorimetric method described by Erel (2004), using dianisidine dihydrochloride as a substrate, at the laboratory of the Biochemistry Department of Atatürk University, Faculty of Veterinary Medicine. The total antioxidant amount of the tarragon plant was determined as 0.833 mmol/g. In another study conducted in the same period and in the same region, the β -ocimene (1 237.21 arbitrary units, AU $\times 10^{-6}$), α -pinene (114.4 AU $\times 10^{-6}$), β -thujene (166.92 AU $\times 10^{-6}$), D-limonene (366.2 AU $\times 10^{-6}$), γ -terpinene (187.27 AU $\times 10^{-6}$), Terpinolene (129.92 AU $\times 10^{-6}$) and Estragole (11 242 AU $\times 10^{-6}$) levels were stated (Yilmaz et al. 2019).

Collection of serum samples

At the end of the trial, 96 laying hens (three birds in each subgroup, a total of twelve birds in each group) were randomly selected for the blood sampling. The samples were collected from the *vena subcutanea ulnaris* to 2 ml tubes during the cervical dislocation. A refrigerated centrifuge (NF 1200, CORE, Ankara, Turkey) at +4 °C for 12 min was used to obtain the serum samples (Tekce and Gul 2015).

Measurement of serum apelin-13 and plasma BDNF concentrations

The serum apelin, and plasma brain-derived neurotrophic factor (BDNF) were measured us-

<https://doi.org/10.17221/9/2020-VETMED>

ing an Enzyme-Linked Immuno Sorbent Assay (ELISA) (R&D Systems, Minneapolis, MN, USA) and the values were read using an ELISA reader (Mindray MR-96 A, P.R. China).

The minimum detectable concentration used to measure the apelin concentration in the blood serum obtained from the research was < 18.75 pg/ml. An ELISA kit type-specific for chicken apelin (FineTest, Product code: ECH0078, P.R. China) from 31.25–2 000.00 pg/ml, an intra-assay coefficient of 8.0%, and an inter-assay coefficient of 10.0% was utilised in accordance with the manufacturer’s protocol.

The results were evaluated by reading the absorbance values at 450 nm in accordance with the procedure reported in the kit (Dai et al. 2018). The minimum detectable concentration to measure serum cTnI concentrations in the blood serum obtained from the study was < 9.4 pg/ml. An ELISA kit type-specific for chicken cTnI (FineTest, Product code: ECH0069, P.R. China) from 15.6–1 000.00 pg/ml, an intra-assay coefficient of 8.0%, and an inter-assay coefficient of 10.0% was utilised in accordance with the manufacturer’s protocol. The results were evaluated by reading absorbances at 450 nm in accordance with the procedure reported in the kit (Bayraktar and Tekce 2019). A commercially available chicken BDNF (Product code: 201-16-1172; Sunred, P.R. China) ELISA kit was utilised by reading the 450 nm absorbance values in accordance with the procedure reported in the kit (Dai et al. 2018).

Histopathologic assessment

At the end of the trial, cervical dislocation was performed in 80 birds in total, randomly selected as ten out of each group, for the histopathological assessment and necropsy. The liver tissue samples taken for the histopathological assessment were then fixed for 48 hours in a 10% formalin solution. They were embedded in paraffin blocks according to the routine histological follow-up procedures. Cross-sections were taken from each block at a 4 µm thickness (Bancroft et al. 2012). The microscopic findings of the hepatic tissue (parenchyma, serosa) samples were assessed for each study group (Leica DM 1000, Germany). In the sub-groups, which were housed at the normal stocking density and received different doses of the dietary tarragon (Groups NT₁, NT_{1,2}, NT₆, NT₁₂), the microscopic examination of the hepatic tissue showed that the histological structure of the parenchyma and serosa was normal. In the sub-groups, housed at the high stocking density, and receiving different doses of the dietary tarragon (Groups ST₁, ST_{1,2}, ST₆, ST₁₂), the histopathological examination of the hepatic tissue samples revealed the hydropic degeneration of the hepatocytes, ranging from very severe, severe, and moderate to mild, and steatosis of the hepatocytes, ranging from very severe and moderate to mild. The sub-groups ST₁, ST_{1,2}, and ST₆ presented with severe hyperaemia of the sinusoidal, portal and acinar blood vessels, whilst in sub-group ST₁₂, the hyperaemia of these blood vessels was moderate (Figure 1).




Figure 1. Histopathological view of the liver tissue (× 40, H&E)

Statistical analysis

The apelin, p-BDNF and cTnI concentrations were controlled for normal distribution and all of them were distributed normally. The statistical analyses of the diet and stocking densities effects on the apelin, p-BDNF and cTnI were performed using the General Linear Model (GLM) that is given below:

$$Y_{ijk} = \mu + D_i + T_j + (D \times T)_{ij} + e_{ijk} \quad (1)$$

where:

- Y_{ijk} – an observation;
- μ – the overall mean;
- D_i – the diet effect;
- T_i – the stocking densities effect;
- $(D \times T)_{ij}$ – the interaction effect;
- e_{ijk} – the experimental error.

A one-way analysis of variance (ANOVA) was used for a 2 (two stocking densities levels) \times 3 (three dose levels) factorial design. The Duncan

multiple comparison test was used for comparing the group means. The statistical analyses were performed by using IBM SPSS statistics v22.0.

RESULTS

In this study, the impact of ground tarragon (*Artemisia dracunculus*), added at different doses (0, 1.2, 6 and 12 g/kg in Groups T₀, T_{1,2}, T₆ and T₁₂) in a homogenous form to the feed of egg-laying hens housed at different stocking densities, was investigated on the apelin, BDNF, and cTnI concentrations. The data pertaining to the study groups are presented in Table 2.

In the group, which was housed at the normal stocking density (810 cm²/hen) and was not exposed to stress, the apelin concentration was the highest in sub-group NT_{1,2} (1.397 ng/ml) and the lowest in sub-group NT₁₂ (0.641 ng/ml). Furthermore, while the p-BDNF concentration was the highest in sub-group NT₆ (2.086 ng/ml) and the lowest in sub-group NT₀ (1.965 ng/ml), the cTnI concentration was the

Table 2. Hormone levels of apelin, BDNF, cTnI due to the tarragon administration in the laying hens found in the stocking density (ng/ml)

	Apelin (ng/ml)		BDNF (ng/ml)		cTnI (ng/ml)	
	non stress	stress	non stress	stress	non stress	stress
T ₀	0.878 ± 0.315	0.874 ± 0.315	1.965 ± 0.031	2.004 ± 0.031	2.110 ± 0.052	1.981 ± 0.052
T _{1,2}	1.397 ± 0.315	1.082 ± 0.315	2.029 ± 0.031	2.025 ± 0.031	2.092 ± 0.052	2.057 ± 0.052
T ₆	1.038 ± 0.315	0.679 ± 0.315	2.086 ± 0.031	2.083 ± 0.031	2.082 ± 0.052	2.084 ± 0.052
T ₁₂	0.641 ± 0.315	1.447 ± 0.315	2.046 ± 0.031	2.009 ± 0.031	1.930 ± 0.052	2.064 ± 0.052
Source of variation (<i>P</i> -values)						
Diet	0.602		0.030		0.363	
Stress	0.883		0.953		0.851	
Stress \times diet	0.249		0.363		0.116	
Main effect means diet						
T ₀	0.878 ± 0.223		1.984 ± 0.022 ^b		2.045 ± 0.037	
T _{1,2}	1.239 ± 0.223		2.027 ± 0.022 ^{ab}		2.074 ± 0.037	
T ₆	0.858 ± 0.223		2.085 ± 0.022 ^a		2.083 ± 0.037	
T ₁₂	1.044 ± 0.223		2.028 ± 0.022 ^{ab}		1.997 ± 0.037	
Stress						
Non stress	0.988		2.032		2.053	
Stress	1.022		2.030		2.046	
SEM	0.158		0.015		0.026	

^{a-c}The means with the same letters in a column are statistically different ($P > 0.05$)

<https://doi.org/10.17221/9/2020-VETMED>

highest in sub-group NT₀ (2.110 ng/ml) and the lowest in sub-group NT₁₂ (1.930 ng/ml).

In the group, which was housed at the high stocking density (580 cm²/hen) and exposed to stress, the apelin concentration was the highest in sub-group ST₁₂ (1.447 ng/ml) and the lowest in sub-group ST₆ (0.679 ng/ml). While the p-BDNF concentration was the highest in sub-group ST₆ (2.083 ng/ml) and the lowest in sub-group ST₀ (2.004 ng/ml), the cTnI concentration was the highest in sub-group ST₆ (2.084 ng/ml) and the lowest in sub-group ST₀ (1.981 ng/ml). The serum apelin concentrations in the groups housed at the normal and high stocking density were determined to fall within the normal reference range, and did not show any statistically significant differences ($P > 0.05$). The analyses demonstrated that the dietary tarragon supplementation had no effect on the apelin, BDNF and cTnI concentrations under stressful or stress-free conditions ($P > 0.05$).

The histopathological findings detected in the study groups are shown in Figure 1. The examinations demonstrated that the groups that received the dietary tarragon under the stress-free conditions did not differ from the control group. On the other hand, when compared to the control group, the study groups that were exposed to the stress presented with very severe hepatocytic steatosis, and the presence of vacuolised lipid degenerations in the cytoplasm of the hepatocytes, which resulted in the peripheral positioning of the nucleus in these cells. The hepatocytes showed very severe hydropic degeneration, their cytoplasm was swollen, and the cells were stained a pale colour. The sinusoidal, portal and acinar blood vessels were severely hyperaemic. When compared to the control group, it was determined that, in the sub-groups, which were exposed to stress and received dietary tarragon, depending on the dose of tarragon, the hepatocyte degeneration, sinusoidal dilatation and hyperaemia of the sinusoidal, portal and acinar blood vessels decreased. Thus, the tarragon was determined to show a regulatory effect.

DISCUSSION

As is the case in several other animal species, in poultry, the body responds to stress with endocrine and biochemical alterations. Increasing the stocking density of birds in cages elevates

the plasma corticosterone concentrations (Eugen et al. 2019), inhibits the locomotor development (Puron et al. 1995; Feddes et al. 2002; Dawkins et al. 2004), and due to the inefficiency of the body heat loss mechanism, results in heat stress, which, in return, reduces the yields and deteriorates the health status of the animals (Onbasilar and Aksoy 2005). While cholesterol (CHOL), glucose (GLU), high-density lipoprotein (HDL), and triglyceride (TRI) concentrations are accepted as stress markers for all poultry species, plasma corticosterone concentrations are not always considered to be an indicator of stress for egg-laying hens (Mumma et al. 2006). In response to the increased metabolic demand, the glucocorticoids mobilise energy reserves, thus, while increased plasma CORT concentrations have positive outcomes in the short-term (Turner et al. 2012), they may cause adverse effects in the long-term (Turner et al. 2010). The data obtained for the plasma CORT concentrations in the present study agree with some literature reports (Fitko et al. 1993; Eugen et al. 2019), but disagree with some other studies (Downing and Bryden 1999; Davis et al. 2000).

Different from mammals, in avian species, lipogenesis mainly occurs in the hepatic tissue and is observed at a very limited level in the adipose tissue (Hermier 1997). Glucocorticoids, released in response to stress, firstly mobilise the adipose tissue. The adipose tissue is the main body source of reactive oxygen species (ROS), which cause oxidative stress when their plasma concentrations are elevated (Furukawa et al. 2004). The hormone apelin, released from the adipose tissue, owing to its interaction with the apelin receptor (APJ), suppresses the production of the reactive oxygen species in the adipocytes. Furthermore, apelin is reported to enhance the expression of the antioxidant enzymes through the activation of the mitogen-activated protein kinase (MAPK)/extracellular signal-regulated protein kinase (ERK) and AMP-activated protein kinase (AMPK) pathways, to inhibit the expression of the pro-oxidant enzyme via the AMPK pathway, and to reduce the oxidative stress-induced disorders of the pro- and oxidant enzymes, mitochondrial biogenesis and functional expression, and the release of proactive and anti-inflammatory adipocytokines (Than et al. 2014). It is accepted that, through the release of various adipocytokines and metabolic factors, the adipose tissue regulates the metabolic homeostasis (Trayhurn

et al. 2006). Research on tarragon, known to have a major impact on adipogenesis, has shown that this plant enhances adipocyte development and positively affects adipocyte-related diseases by increasing the level of apelin and adiponectin released from the adipose tissue, and strengthening the effect of the insulin hormone (Richard et al. 2014). Moreover, it has been reported that tarragon shows an effect on the membrane of the skeletal cells, increases the insulin sensitivity and adipocyte differentiation in cultures (Anaya-Eugenio et al. 2014; Obanda et al. 2014; Richard et al. 2014). In a similar study conducted in rats, it was determined that the extract of *Artemisia dracuncululus* leaves could potentially decrease the prevalence of coronary diseases in humans (Yazdanparast and Saei 1999; Duric et al. 2015). Stress causes damage to the hepatic parenchyma and sinusoidal structures. Perfusion and the generation of free radicals cause neutrophil leukocyte infiltration in the liver and hyperaemia in tissues. In the present study, it was ascertained that a stocking density not causing any stress to animals had no adverse effect on the liver, and produced results similar to those of the control group, as has also been suggested in previous research (Shen et al. 2007). On the other hand, when compared to the control group, in the sub-groups exposed to stress, very severe steatosis, the presence of vacuolised lipid degeneration, cytoplasmic lipid vacuoles and peripheral nuclear dislocation were observed in the hepatocytes, and the severity of these findings was observed to decrease with the increased doses of the dietary tarragon. This was attributed to the strong antioxidant property of the tarragon plant (*Artemisia dracuncululus*).

When chicken flocks are housed at a high stocking density, the dominant animals take hold of first access to the feed, water and other valuable resources (Estevez 2002). Stress is known to cause arrhythmia in the ventricular system of the heart (Scorza et al. 2010). In recent years, cardiac troponin I (cTnI) has found common use as a highly specific cardiac marker for stress-induced cardiac arrhythmia, myocardial damage and myocardial disease (Adams et al. 1994; Wu et al. 1996; Brown and Bertolet 1997). Apelin-13 has a role in regulating the stress response with the BDNF by improving the HPA Axis and Hippocampal Glucocorticoid Receptor Dysfunctions (Dai et al. 2018). Apelin has a protective effect against oxidative stress in heart myocardial cells as in many tissues (Chung et al.

2016). Besides, the hormones apelin and BDNF also have regulatory roles in cardiac contraction (Szokodi et al. 2002; Fulgenzi et al. 2015). On the other hand, BDNF, one of the key adipokines involved in neurodegenerative processes, shows a neuroprotective effect (Dai et al. 2018) and acts as the direct modulator of myocardial mechanic function in the BDNF/TrkB signalling (Feng et al. 2015). To the authors' knowledge, the impact of dietary tarragon on the serum apelin levels in egg-laying hens housed at different stocking densities has not been investigated before. The results obtained in the present study for the apelin, p-BDNF, and cTnI levels are similar to those indicated in some literature reports (Feng et al. 2015; Dai et al. 2018; Shen et al. 2019). As the apelin, p-BDNF, and cTnI levels have not been investigated in poultry before, the data of the present study was compared to the results of previous research conducted in different species.

In conclusion, dietary tarragon supplementation had no effect on the apelin, BDNF and cTnI concentrations under stressful or stress-free conditions ($P > 0.05$). However, based on the results of this study, which is the first investigation on the apelin, p-BDNF, and cTnI concentrations in laying hens raised at different stocking densities, it is considered that these indicators can be used as hormonal markers of the response to stress, and thus, can aid in the prediction of the stress, in the assessment of the animal's welfare and in the development of innovative livestock management strategies. Furthermore, more detailed research is needed in this area to confirm the results of this study.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Adlard PA, Cotman CW. Voluntary exercise protects against stress-induced decreases in brain-derived neurotrophic factor protein expression. *Neuroscience*. 2004 Dec 23;124(4):985-92.
- Adams JE, Sicard GA, Allen BT, Bridwell KH, Lenke LG, Davila-Román VG, Bodor GS, Ladenson JH, Jaffe AS. Diagnosis of perioperative myocardial infarction with measurement of cardiac troponin I. *N Engl J Med*. 1994 Mar 10;330(10):670-4.

<https://doi.org/10.17221/9/2020-VETMED>

- Aglarova AM, Zilfikarov IN, Severtseva OV. Biological characteristics and useful properties of tarragon (*Artemisia dracunculus* L.). *Pharm Chem*. 2008 Aug 10;42(2):81-6.
- Alpert JS, Thygesen K, Antman E, Bassand JP. Myocardial infarction redefined – A consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *J Am Coll Cardiol*. 2000 Sep;36(3):959-69.
- Anaya-Eugenio GD, Rivero-Cruz I, Rivera-Chavez J, Mata R. Hypoglycemic properties of some preparations and compounds from *Artemisia ludoviciana*. *Nutt J Ethnopharmacol*. 2014 Aug 8;155(1):416-25.
- Apple FS. Clinical and analytical standardization issues confronting cardiac troponin I. *Clin Chem*. 1999 Jan 1;45(1):18-20.
- Appleby MC. What causes crowding? Effects of space, facilities and group size on behaviour, with particular reference to furnished cages for hens. *Anim Welf*. 2004 Aug 3;13(3):313-20.
- Appleby MC, Smith SE, Hughes BO. Nesting, dust bathing and perching by laying hens in cages: Effects of design on behaviour and welfare. *Br Poult Sci*. 1993 Apr 27;34(5):835-47.
- Attokaran M. Natural food flavors and colorants. Oxford: Blackwell Publishing Ltd. and Institute of Food Technologists; 2011 Feb 7. p. 377-8.
- Bayraktar B, Tekce E. Effects of varying essential oil mixture concentrations applied under conditions of different temperature stress on cardiac markers and other blood parameters. *Braz J Poultry Sci*. 2019 Dec 20;21(4):1-8.
- Bancroft JD, Suvarna K, Layton C. Bancroft's theory and practice of histological techniques. 7th ed. Philadelphia, PA, USA: Churchill Livingstone, Elsevier; 2012. 654 p.
- Beltowski J. Apelin and visfatin: Unique “beneficial” adipokines upregulated in obesity? *Med Sci Monit*. 2006 Jun;12(6):RA112-9.
- Benraiss A, Chmielnicki E, Lerner K, Roh D, Goldman SA. Adenoviral brain-derived neurotrophic factor induces both neostriatal and olfactory neuronal recruitment from endogenous progenitor cells in the adult forebrain. *J Neurosci*. 2001 Sep 1;21(17):6718-31.
- Bessei W. Welfare of broilers: A review. *Worlds Poult Sci J*. 2006 Sep 1;62(3):455-66.
- Bothwell M. Functional interactions of neurotrophins and neurotrophin receptors. *Annu Rev Neurosci*. 1995;18:223-53.
- Brown CS, Bertolet BD. Cardiac troponin: See ya later, CK! *Chest*. 1997 Jan 1;111(1):2-4.
- Ceylan A. *Tıbbi Bitkiler II (Uçucu Yağ Bitkileri) [Medicinal Plants II (Essential Oil Plants)]*. İzmir: Ege University Faculty of Agriculture; 1996. 481 p. Turkish.
- Charmandari E, Tsigos C, Chrousos G. Endocrinology of the stress response. *Annu Rev Physiol*. 2005;67:259-84.
- Chung WJ, Cho A, Byun K, Moon J, Ge X, Seo HS, Moon E, Dash R, Yang PC. Apelin-13 infusion salvages the peri-infarct region to preserve cardiac function after severe myocardial injury. *Int J Cardiol*. 2016 Nov 1;222:361-7.
- Craig JV, Swanson JC. Welfare perspectives on hens kept for egg production. *Poultry Sci*. 1994 Mar 15;73(7):921-38.
- Dai TT, Wang B, Xiao ZY, You Y, Tian SW. Apelin-13 up-regulates BDNF against chronic stress-induced depression-like phenotypes by ameliorating HPA axis and hippocampal glucocorticoid receptor dysfunctions. *Neuroscience*. 2018 Oct 15;390:151-9.
- Davis GS, Anderson KE, Carroll AS. The effects of long-term caging and molt of Single Comb White Leghorn hens on heterophil to lymphocyte ratios, corticosterone and thyroid hormones. *Poultry Sci*. 2000 Apr 1;79(4):514-8.
- Dawkins MS, Donnelly CA, Jones TA. Chicken welfare is influenced more by housing conditions than by stocking density. *Nature*. 2004 Jun 22;427(6972):342.
- Downing JA, Bryden WL. Stress, hen husbandry and welfare: Literature Review of Stress in Poultry. Australia: RIRDC/EIRDC Publication; 1999. 58 p.
- Duke JA. Handbook of medicinal herbs. 2nd ed. Boca Raton, FL: CRC Press; 2002. 870 p.
- Duric K, Kovac Besovic EE, Niksic H, Muratovic S, Sofic E. Anticoagulant activity of some *Artemisia dracunculus* leaf extracts. *Bosn J Basic Med Sci*. 2015 May 13;15(2):9-14.
- Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin Biochem*. 2004 Nov 28;37(4):277-85.
- Estevez I. Density allowances for broilers: Where to set the limits? *Poult Sci*. 2007 Jun;86(6):1265-72.
- Eugen KV, Nordquist RE, Zeinstra E, Staay FJV. Stocking density affects stress and anxious behavior in the laying hen chick during rearing. *Animals (Basel)*. 2019 Feb 10;9(2):53.
- Feddes JJ, Emmanuel EJ, Zuidhof MJ. Broiler performance, body weight variance, feed and water intake, and carcass quality at different stocking densities. *Poultry Sci*. 2002 Jun 1;81(6):774-9.
- Feng N, Huke S, Zhu G, Tocchetti CG, Shi S, Aiba T, Kaludercic N, Hoover DB, Beck SE, Mankowski JL, Tomaselli GF, Bers DM, Kass DA, Paolucci N. Constitutive BDNF/TrkB signaling is required for normal cardiac contraction and relaxation. *Proc Natl Acad Sci USA*. 2015 Feb 10;112(6):1880-5.
- Fitko R, Jakubowski K, Zielinski H, Potrzuska I. Słезenie korlykosleronu, odrenoliny i norodrenoliny oraz oklywosc gronuloocytow obolefnoclonnych we krwi kurczql w

<https://doi.org/10.17221/9/2020-VETMED>

- oslrym slesie immobilizacji [The level of corticosterone, adrenaline and noradrenaline and activity of blood neutrophils in chickens under an acute immobilization stress]. *Med Weter.* 1993;49:37-8. Polish.
- Fuchikami M, Morinobu S, Kurata A, Yamamoto S, Yamawaki S. Single immobilization stress differentially alters the expression profile of transcripts of the brain-derived neurotrophic factor (BDNF) gene and histone acetylation at its promoters in the rat hippocampus. *Int J Neuropsychoph.* 2009 Feb 1;12(1):73-82.
- Fulgenzi G, Tomassoni-Ardori F, Babini L, Becker J, Barrick C, Puverel S, Tessarollo L. BDNF modulates heart contraction force and long-term homeostasis through truncated TrkB.T1 receptor activation. *J Cell Biol.* 2015 Sep 14;210(6):1003-12.
- Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest.* 2004 Dec 15;114(12):1752-61.
- Gulpinar Y. Tarhun Bitkisinin (*Artemisia dracunculus* L.) Wistar Albino Ratlarda Oluşturulmuş Akut Karaciğer Toksik Hasarına Karşı Koruyucu ve Tedavi Edici Etkisinin Araştırılması [Tarragon Plant (*Artemisia dracunculus* L.) Acute Liver Created in Wistar Albino Rats Protective and Therapeutic Effect Against Toxic Damage] [masters' thesis]. Gaziantep: Gaziantep University Science Institute of Biology, Master's Thesis; 2012. Turkish.
- Hassanzadeh MK, Najaran ZT, Nasery M, Emami SA. Tarragon (*Artemisia dracunculus* L.) oils. In: Preedy VR, editor. *Essential oils in food preservation, flavor and safety.* Amsterdam: Elsevier Academic Press; 2016. p. 813-7.
- Hermier D. Lipoprotein metabolism and fattening in poultry. *J Nutr.* 1997 May 5;127(5):805S-8S.
- Hi W, Ta S, An AB. Breast muscle characteristics of avian pathogenic *Escherichia coli* infected broilers fed with antibiotics or probiotic. *Poult Sci J.* 2019 Oct 22;7(2):131-40.
- Hosseinzadeh Z, Moghaddam GA. Effects of tarragon powders' different levels (*Artemisia dracunculus*) on general performance and anetometric properties of digestive system of male broiler chickens. *Int J Adv Biol Biom Res.* 2014;2(5):1599-605.
- Kang HK, Park SB, Kim SH, Kim CH. Effects of stock density on the laying performance, blood parameter, corticosterone, litter quality, gas emission and bone mineral density of laying hens in floor pens. *Poult Sci.* 2016 Jun 15;95(12):2764-70.
- Kertes DA, Bhatt SS, Kamin HS, Hughes DA, Rodney NC, Mulligan CJ. BDNF methylation in mothers and newborns is associated with maternal exposure to war trauma. *Clin Epigenetics.* 2017 Jun 30;9(1):68.
- Kordali S, Kotan R, Mavi A, Cakir A, Ala A, Yildirim A. Determination of the chemical composition and antioxidant activity of the essential oil of *Artemisia dracunculus* and of the antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *A. dracunculus*, *Artemisia santonicum*, and *Artemisia spicigera* essential oils. *J Agric Food Chem.* 2005 Oct 29;53(24):9452-8.
- Lee T, Saruta J, Sasaguri K, Sato S, Tsukinoki K. Allowing animals to bite reverses the effects of immobilization stress on hippocampal neurotrophin expression. *Brain Res.* 2008 Feb 21;1195:43-9.
- Mansuroglu S, Gurel E. Mikroçoğaltım, Bitki Biyoteknolojisi I, Doku Kültürü ve Uygulamaları [Micropropagation, plant biotechnology I: Tissue culture and applications]. Babaoglu M, Gurel E, Ozcan S, editors. Konya: Selcuk University Foundation Publications; 2001. p. 262-81.
- McEwen BS. Protective and damaging effects of stress mediators: Central role of the brain. *Dialogues Clin Neurosci.* 2006 Dec 8;(4):367-81.
- McGlone J. Guide for the care and use of agricultural animals in research and teaching. 3rd ed. Champaign, IL: Federation of Animal Science Societies; 2010. p. 111-2.
- Mumma JO, Thaxton JP, Vizzier-Thaxton Y, Dodson WL. Physiological stress in laying hens. *Poult Sci.* 2006;85(4):761-9.
- Obanda DN, Ribnicky DM, Raskin I, Cefalu WT. Bioactives of *Artemisia dracunculus* L. enhance insulin sensitivity by modulation of ceramide metabolism in rat skeletal muscle cells. *Nutrition.* 2014 Jul-Aug;30(7-8):S59-S66.
- Onbasilar EE, Aksoy FT. Stress parameters and immune response of layers under different cage floor and density conditions. *Livest Prod Sci.* 2005 Aug;95(3):255-63.
- Matkovic K, Marusic D, Ostovic M, Pavicic Z, Matkovic S, Kabalin AE, Lucic H. Effect of litter type and perches on footpad dermatitis and hock burn in broilers housed at different stocking densities. *S Afr J Anim Sci.* 2019 Jun 19;49(3):546-54.
- Meseret S. A review of poultry welfare in conventional production system. *Livest Res Rural Dev.* 2016 Dec 1;28:12-8.
- Nicol CJ, Bouwsema J, Caplen G, Davies AC, Hockenhill J, Lambton SL, Mullan S, Weeks CA. Farmed bird welfare science review. Melbourne: Department of Economic Development, Jobs, Transport and Resources; 2017. p. 17-8.
- Puron D, Santamaria R, Segura JC, Alamilla JL. Broiler performance at different stocking densities. *J Appl Poultry Res.* 1995 Mar;4(1):55-60.
- Richard AJ, Burriss TP, Sanchez-Infantes D, Wang Y, Ribnicky DM, Stephens JM. *Artemisia* extracts activate PPAR γ , promote adipogenesis, and enhance insulin sensitivity in adipose tissue of obese mice. *Nutrition.* 2014 Jul-Aug;30(7-8):S31-S6.

<https://doi.org/10.17221/9/2020-VETMED>

- Rogers J, Kochunov P, Zilles K, Shelledy W, Lancaster J, Thompson P, Duggirala R, Blangero J, Fox PT, Glahn DC. On the genetic architecture of cortical folding and brain volume in primates. *Neuroimage*. 2010 Nov 15;53(3):1103-8.
- Scorza FA, Albuquerque RD, Arida RM, Albuquerque MD, Terra VC, Machado HR, Cysneiros RM, Scorza CA, Cavalheiro EA. What are the similarities between stress, sudden cardiac death in *Gallus gallus* and sudden unexpected death in people with epilepsy. *Arq Neuropsiquiatr*. 2010 Oct;68(5):788-90.
- Shen L, Kondo Y, Ahmed S, Bumber Y, Konishi K, Guo Y, Chen X, Vilaythong JN, Issa JP. Drug sensitivity prediction by CpG island methylation profile in the NCI-60 cancer cell line panel. *Cancer Res*. 2007 Dec 1;67(23):11335-43.
- Shen P, Yue Q, Fu W, Tian SW, You Y. Apelin-13 ameliorates chronic water-immersion restraint stress-induced memory performance deficit through upregulation of BDNF in rats. *Neurosci Lett*. 2019 Mar 23;696:151-5.
- Sorensen P, Su G, Kestin SC. Effects of age and stocking density on leg weakness in broiler chickens. *Poult Sci*. 2000 Jun 1;79(6):864-70.
- Spina MB, Squinto SP, Miller J, Lindsay RM, Hyman C. Brain-derived neurotrophic factor protects dopamine neurons against 6-hydroxydopamine and N-methyl-4-phenylpyridinium ion toxicity: Involvement of the glutathione system. *J Neurochem*. 1992 Jul;59(1):99-106.
- Szokodi I, Tavi P, Foldes G, Voutilainen-Myllylä S, Ilves M, Tokola H, Pikkarainen S, Piihola J, Rysä J, Tóth M, Ruskoaho H. Apelin, the novel endogenous ligand of the orphan receptor APJ, regulates cardiac contractility. *Circ Res*. 2002 Sep 6;91(5):434-40.
- Tatemoto K, Hosoya M, Habata Y, Fujii R, Kakegawa T, Zou MX, Kawamata Y, Fukusumi S, Hinuma S, Kitada C, Kurokawa T, Onda H, Fujino M. Isolation and characterization of a novel endogenous peptide ligand for the human APJ receptor. *Biochem Biophys Res Commun*. 1998 Oct 20;251(2):471-6.
- Tatemoto K, Takayama K, Zou MX, Kumaki I, Zhang W, Kumano K, Fujimiya M. The novel peptide apelin lowers blood pressure via a nitric oxide-dependent mechanism. *Regul Pept*. 2001 Jun 15;99(2-3):87-92.
- Tekce E, Gul M. Sicaklık Stresi Altında Beslenen Etçi Piliçlerde *Origanum Syriacum* Uçucu Yağının performans Antioksidan Potansiyel Lipid Profili Bağırsak Mikrobiyotası ve Et Kalitesine Etkisi [dissertation]. AÜ Sağlık Bilimleri Enstitüsü, Erzurum. 2015. Turkish.
- Than A, Zhang X, Leow MK, Poh CL, Chong SK, Chen P. Apelin attenuates oxidative stress in human adipocytes. *J Biol Chem*. 2014 Feb 7;289(6):3763-74.
- Trayhurn P, Bing C, Wood IS. Adipose tissue and adipokines – Energy regulation from the human perspective. *J Nutr*. 2006 Jul;136(7 Suppl):1935S-9S.
- Turner AI, Keating CL, Tilbrook AJ. Sex differences and the role of sex steroids in sympatho-adrenal medullary system and hypothalamo-pituitary-adrenal axis responses to stress. *Sex steroids*. 2012 Jan;12:115-36.
- Turner AI, Rivalland ET, Clarke IJ, Tilbrook AJ. Stressor specificity of sex differences in hypothalamo-pituitary-adrenal axis activity: Cortisol responses to exercise, endotoxin, wetting, and isolation/restraint stress in gonadectomized male and female sheep. *Endocrinology*. 2010 Sep 1;151(9):4324-31.
- Voitkevich SA. [Essential oils for perfumery and aromatherapy]. Moscow: Pishchevaya Promyshlennost'; 1999. p. 212-3. Russian.
- Volak J, Stodola J. The illustrated book of herbs. London: Caxton Editions; 1998. 256 p.
- Walzem RL, Chen SE. Obesity-induced dysfunctions in female reproduction: Lessons from birds and mammals. *Adv Nutr*. 2014 Mar 1;5(2):199-206.
- Weimer SL, Robison CI, Tempelman RJ, Jones DR, Karcher DM. Laying hen production and welfare in enriched colony cages at different stocking densities. *Poult Sci*. 2019 Sep 1; 98(9):3578-86.
- Wu AH, Feng YJ, Contois JH, Pervaiz S. Comparison of myoglobin, creatine kinase-MB, and cardiac troponin I for diagnosis of acute myocardial infarction. *Ann Clin Lab Sci*. 1996 Jul 1;26(4):291-300.
- Yazdanparast R, Saei A. Effects of aqueous tarragon, *Artemisia dracunculus*, extract on lipid and coagulatory parameters in rats. *Biomed Lett*. 1999;59(233):137-41.
- Yilmaz O, Kaban G, Kaya M. Essential oil compounds of tarragon and coriander seed. *Bayburt University J Sci*. 2019 May 21;2(1):17-24.
- Zigova T, Pencea V, Wiegand SJ, Luskin MB. Intraventricular administration of BDNF increases the number of newly generated neurons in the adult olfactory bulb. *Mol Cell Neurosci*. 1998 Jul;11(4):234-45.

Received: January 1, 2020

Accepted: May 28, 2020